

Commentary

It Is Not What We Eat, It's How We Produce It

Jaime Martínez-Valderrama 

Estación Experimental de Zonas Áridas, CSIC, La Cañada de San Urbano, Almería 04120, Spain; jaimonides@eeza.csic.es

1. Exceeding Critical Thresholds in Food Production

The current food system is one of the main contributors to the environmental degradation of our planet (FAO, 2021). The disruption of phosphorus and nitrogen cycles, the release of greenhouse gases (GHGs), and the degradation of vital resources like water and soil serve as compelling examples and explain the overtaking of six planet boundaries (Rockström & Gupta, 2023) that a decade ago appeared distant and abstract.

Despite this prevailing consensus, there is a substantial divergence in the range of measures and proposals. Many focus on product analyses, emphasizing their water and carbon footprint (Mekonnen & Hoekstra, 2012; Petterson et al., 2021), as well as their relationship with health (Willett et al., 2019) often overlook the significance of production systems and our lifestyle. In an era marked by short-termism and the incessant flow of information, short and flashy messages are imposed, leaving aside essential nuances of the food system, such as the externalities of each form of production, the demand for very cheap food at any time, or the final form in which these products are delivered to us (ultra-processed, with a huge associated consumption of plastic and energy, or with working conditions that are in many cases appalling).

It is imperative to expand our perspective and take into account the diverse impacts—both positive and negative—across social, environmental, and economic dimensions of food systems. This is crucial as the same product can be cultivated and produced through various methods, each carrying distinct implications.

This viewpoint can be illustrated with two examples. One concerns the call to drastically reduce red meat consumption, and the other promotes so-called superfoods (Magrach & Sanz, 2020), such as quinoa. Initially, several data points indeed underpin the sustainability of this dietary shift. Beef production accounts for an average water consumption of 15,415 m³ t⁻¹ (Mekonnen & Hoekstra, 2012). 98% of this water footprint is due to the massive use of animal feed, the production of which requires the cultivation of cereals and legumes. The cultivation of these crops involves the deforestation of valuable primary forests (Martínez-Valderrama et al., 2021) and the depletion of aquifers, which are also affected by the discharge of animal slurry from large-scale farms. Another harmful effect associated with livestock farming is the emission of GHGs, mainly methane and nitrous oxide. It is estimated that this production sector is responsible for between 8 and 18% of total emissions (Herrero & Thornton, 2013); on average, meat carbon footprint is 41 kg CO₂eq kg⁻¹ (Herrero et al., 2013). In addition, there are other undesirable effects of meat production and consumption, such as the conditions in which many of these animals live and the diseases associated with excessive consumption. Quinoa enjoys a favorable reputation. It's a food with deep roots in ancient cultures, which imparts a sense of exoticism in Western markets. Often hailed as a “superfood”, quinoa is highly compatible with vegetarian diets, offering a rich blend of essential amino acids, micronutrients, vitamins, and is naturally gluten-free.

2. The Need for a More Integrated Vision of Food Systems

Is this always the case? It depends, we can argue the opposite. Let us first look at meat. There are many livestock production systems that are examples of sustainability, i.e., stocking rates are adjusted to the availability of pasture. Pastoralism is the most widespread land use around the world and has proven to be a secure livelihood for many societies for millennia (Manzano et al., 2021). In these livestock grazing systems, the ruminants graze in the open air, eating various types of vegetation that would otherwise not be utilized. In doing so, they achieve something unique: they convert lignin and cellulose into protein. No machine is capable of this process. In addition, they rid the landscape of flammable materials, reducing the risk of forest fires. As they move, they fertilize the countryside with their excrement, in line with the precepts of the circular economy: that the waste of one becomes the food of the other. The richness of breeds is the result of their adaptability to different environments and conditions, which translates into great agrobiodiversity and resilience of the territory. Finally, products derived from this type of livestock farming have

Citation: Martínez-Valderrama, J. It is Not What We Eat, It's How We Produce it.

Agricultural & Rural Studies, 2024, 2, 0006.

<https://doi.org/10.59978/ar02010006>

Received: 6 November 2023

Revised: 20 November 2023

Accepted: 29 November 2023

Published: 7 February 2024

Publisher's Note: Trenton Gary, SCC Press stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2024 by the author(s). Licensee Trenton Gary, SCC Press, Kowloon, Hong Kong S.A.R., China. The article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

nutritional and organoleptic properties far superior to those of processed products from industrial livestock farming (Wilkinson & Lee, 2018).

What are the downsides of quinoa's popularity? Its soaring demand has pushed it into large supermarket chains where competitive pricing is the standard. As a result, the traditional Andean production system has undergone significant alterations. Peru and Bolivia, the primary quinoa producers, have increased their production by 252% and 612%, respectively, over the last four decades, along with a 124% and 440% expansion in their cultivation areas (Magrach & Sanz, 2020). These changes have brought about noteworthy environmental and social consequences. In the high-altitude deserts where quinoa thrives, the conditions are harsh: minimal rainfall coupled with frigid and windy weather. For centuries, local communities sustained themselves on quinoa, cultivating it without the use of mechanical equipment and meticulously considering weather conditions to allow the soil to rest and regenerate with water and nutrients. The quinoa boom, which drove prices to unprecedented levels (sometimes reaching up to 60 times that of wheat), triggered the intensification and expansion of quinoa farming.

The use of heavy machinery, fertilizers and pesticides, the elimination of livestock that fertilized the soil, the invasion of pastures and the reduction of fallow have triggered soil erosion and deterioration. The selection of the most productive varieties (4 of which account for 90% of production) is leading to the loss of a rich gene bank. The local population, far from getting richer from this business, has lost its main sources of protein: Quinoa has prohibitive prices and most of it is exported -Peru went from exporting 60 t in 1995 to 36,000 in 2014 (Bedoya-Perales et al., 2018) and the llamas have less space to graze. To make matters worse, much of the land that was in the hands of local communities is now private property.

As evident, a plate of quinoa can have an environmental impact comparable to that of a hamburger. Concentrating solely on the product type, neglecting the social and environmental repercussions of its production system, may result in the formulation of policies that exacerbate rather than resolve issues. Therefore, adopting a more holistic perspective on food systems can empower stakeholders to devise more sustainable land-use plans.

3. Final Remarks

These two instances illustrate that consumption should be guided not solely by the type of food but also by the methods of its production. The main problem is related to large-scale production, which seeks to minimize production costs at the expense of social and environmental externalities (Martínez-Valderrama et al., 2023). This phenomenon of "Uberization" has permeated a significant portion of food systems, and this is where attention should be directed.

Quinoa can maintain its sustainability if cultivation respects local ecosystems and traditional production systems, such as fallowing, and refrains from encroaching upon marginal areas traditionally designated for grazing. On the other hand, livestock production requires substantial reforms, which may include the following guidelines: (i) By reducing animal protein demand for the nutritional reasons outlined above; (ii) By favoring pasture-based livestock systems, within the limits of adequate stocking rates; or (iii) by further technifying intensive production systems (i.e., macro-farms) through cultured meat and precision fermentation (Singh et al., 2022), which will reduce the environmental footprint and animal suffering.

Finally, it is necessary to understand that the elimination of negative externalities and the establishment of socially equitable food systems will result in higher food costs (Baker et al., 2020). The impact on society can be mitigated by redistributing these costs or by narrowing the profit margins of major distributors.

CRedit Author Statement: This is a single author paper, and the author was solely responsible for the content, including the concept, analysis, writing, and revision of the manuscript.

Data Availability Statement: Not applicable.

Funding: This research is supported by ATLAS project, funded by the Biodiversity Foundation of the Ministry for the Ecological Transition and the Demographic Challenge (MITECO) within the framework of the Recovery, Transformation and Resilience Plan (PRTR), financed by the European Union - NextGenerationEU.

Conflicts of Interest: The author declares no conflict of interest.

Acknowledgments: Not applicable.

References

- Bedoya-Perales, N. S., Pumi, G., Talamini, E. & Padula, A. D. (2018). The quinoa boom in Peru: Will land competition threaten sustainability in one of the cradles of agriculture? *Land Use Policy*, 79, 475–480. <https://doi.org/10.1016/j.landusepol.2018.08.039>
- Baker, L., Castilleja, G., De Groot Ruiz, A., Jones, A. (2020). Prospects for the True Cost Accounting of Food Systems. *Nat. Food*, 1, 765–767. <https://doi.org/10.1038/s43016-020-00193-6>

- Food and Agriculture Organization of the United Nations. (2021) *The state of the world's land and water resources for food and agriculture. Systems at breaking point*. Synthesis report. <https://www.fao.org/3/cb7654en/cb7654en.pdf>
- Herrero, M., Havlik, P., Valin, H., Notenbaert, A., Rufino, M. C., Thornton, P. K., Blümmel, M., Weiss, F., Grace, D. & Obersteiner, M. (2013). Biomass use, production, feed efficiencies, and greenhouse gas emissions from global livestock systems. *Proceedings of the National Academy of Sciences*, 110(52), 20888–20893.
- Herrero, M. & Thornton, P. K. (2013). Livestock and global change: Emerging issues for sustainable food systems. *Proceedings of the National Academy of Sciences*, 110(52), 20878–20881. <https://doi.org/10.1073/pnas.1321844111>.
- Magrach, A. & Sanz, M. J. (2020). Environmental and social consequences of the increase in the demand for 'superfoods' world-wide. *People and Nature*, 2(2), 267–278. <https://doi.org/https://doi.org/10.1002/pan3.10085>
- Manzano, P., Burgas, D., Cadahía, L., Eronen, J. T., Fernández-Llamazares, Á., Bencherif, S., Holand, Ø., Seitsonen, O., Byambaa, B., Fortelius, M., Fernández-Giménez, M. E., Galvin, K. A., Cabeza, M. & Stenseth, N. C. (2021). Toward a holistic understanding of pastoralism. *One Earth*, 4(5) 651–665. <https://doi.org/10.1016/j.oneear.2021.04.012>
- Martínez-Valderrama, J., Sanjuán, M. E., del Barrio, G., Guirado, E., Ruiz, A. & Maestre, F. T. (2021). Mediterranean landscape re-greening at the expense of South American agricultural expansion. *Land*, 10(2), 204. <https://doi.org/10.3390/land10020204>
- Martínez-Valderrama, J., Olcina, J., Gartzia, R., Guirado, E., Ibáñez, J. & Maestre, F. T. (2023). Uberizing agriculture in drylands: A few enriched, everyone endangered. *Water Resources Management*, 1–22. <https://doi.org/10.1007/s11269-023-03663-1>
- Mekonnen, M. M., & Hoekstra, A. Y. (2012). A global assessment of the water footprint of farm animal products. *Ecosystems*, 15(3), 401–415. <https://doi.org/10.1007/s10021-011-9517-8>
- Petersson, T., Secondi, L., Magnani, A., Antonelli, M., Dembska, K., Valentini, R., Varotto, A. & Castaldi, S. A. (2021). Multilevel carbon and water footprint dataset of food commodities. *Scientific Data*, 8(1), 127. <https://doi.org/10.1038/s41597-021-00909-8>
- Rockström, J., & Gupta, J. (2023). Safe and just Earth system boundaries. *Nature*, 619, 102–111. <https://doi.org/10.1038/s41586-023-06083-8>
- Singh, S., Yap, W. S., Ge, X. Y., Min, V. L. X. & Choudhury, D. (2022). Cultured meat production fuelled by fermentation. *Trends in Food Science & Technology*, 120, 48–58. <https://doi.org/10.1016/j.tifs.2021.12.028>
- Wilkinson, J. M., & Lee, M. R. F. (2018). Review: Use of human-edible animal feeds by ruminant livestock. *Animal*, 12, 1735–1743. <https://doi.org/10.1017/S175173111700218X>
- Willett, W., Rockström, J., Loken, B., Springmann, M., Lang, T., Vermeulen, S., Garnett, T., Tilman, D., DeClerck, F., Wood, A., Jonell, M., Clark, M., Gordon, J. L., Fanzo, J., Hawkes, C., Zurayk, R., Rivera, J. A., Vries, W. D., Sibanda, L. M., ... & Murray, C. J. L. (2019). Food in the Anthropocene: The EAT–Lancet Commission on healthy diets from sustainable food systems. *The Lancet*, 393(10170), 447–492. [https://doi.org/10.1016/S0140-6736\(18\)31788-4](https://doi.org/10.1016/S0140-6736(18)31788-4)